Design For Your Health

Kukje Pharma's Product List

KUKJE PHARMA

· www.kukjepharm.co.kr · trading@kukjepharm.co.kr

Head Office: 96-8, Yatap-ro, Bundang-gu, Seongnam-si, Gyeonggi-do, Republic of Korea Tel: 82.31.781.9081 / Fax: 82.31.781.6043 / Email: trading@kukjepharm.co.kr
Plant: 47, Sanseong-ro, Danwon-gu, Ansan-si, Gyeonggi-do, Republic of Korea Tel: 82.31.491.9411 / Fax: 82.31.492.5177

2509010F-FX

KUKJE PHARMA

Your Reliable Partner

KUKJE PHARMA

Contents

Ophthalmic Agents	01-05
Cephalosporins	05-07
Penicillines Quinolones Tetracyclines Macrolides	07-09
Antifungal & Antiviral Agents Adrenal Corticosteroids	09-10
CNS Agents Osteoporotic Agents Antihistamines	10-12
NSAIDs & Other Analgesics Gastrointestinal Agents	12-14
Cardiovascular Agents Antidiabetics	15-19
Genitourinary Agents Respiratory Agents	19-21
Other ETCs	21
Health Functional Food	21

Ophthalmic Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Diabetic retinopathy preparations **Tagen F** soft cap. **Tagen F** tab.		Each capsule contains, Bilberry fruit dried Ex. 170mg(60mg as Anthocyanoside)	· Retinopathy, nyctalopia · 1 cap 2~3 times/day
Diabetic retinopathy preparatons **Retium** tab. **250mg, 500mg**		Each tablet contains, Dobesilate calcium Hydrate 250mg, 500mg	· Vascular damage with increased capillary rupture and permeability disorder, diabetic capillary dysfunction, diabetic retinopathy, venous dysfunction, post-thrombotic syndrome, peripheral congestive edema, hemorrhoids. · Vascular damage and retinopathy : 500~1000mg daily · Venous disease : 750mg daily for 1 to 3weeks, then 500mg daily
Dry eye preparations **RebaEye** **ophthalmic solution 2%** **5mL**		Each 1mL contains, Rebamipide 20mg	· Improvement of corneal conjunctival epithelial disorder in adult dry eye syndrom patient · 1 drop 4 times daily
Dry eye preparations **RebaEye** **ophthalmic solution 2%** **0.4mL**		Each 1mL contains, Rebamipide 20mg	· Improvement of corneal conjunctival epithelial disorder in adult dry eye syndrom patient · 1 drop 4 times daily
Dry eye preparations **Cualone** eye drops **(multi dose)** **0.1%** **5mL, 10mL**		Each 1mL contains, Sodium hyaluronate 1mg	· Keratoconjunctival epithelial damage (caused by sjogren syndrome, dry eye syndrome, trauma, hard contact lens, etc.) · 1 drop 5~6 times daily
Dry eye preparations **Cualone** eye drops **(single dose)** **0.1% - 0.45mL** **0.15% - 0.5mL** **0.18% - 0.45mL** **0.3% - 0.5mL**		Each 1mL contains, Sodium hyaluronate 1mg, 1.5mg, 1.8mg, 3mg	· Keratoconjunctival epithelial damage (caused by sjogren syndrome, dry eye syndrome, trauma, hard contact lens, etc.) · 1 drop 5~6 times daily · Use immediately after opening and discard remaining fluids and containers
Dry eye preparations **Restaporin** eye drops **0.05%** **0.4mL**		Each 1mL contains, Cyclosporine 0.5mg	· 1 drop twice a day in each eye approximately 12 hours apart.
Dry eye preparations **Diqueye** eye drops 3% **0.4mL**		Each 1mL contains, Diquafosol sodium 30mg	· Treatment of the signs and symptoms of dry eye disease · One drop 6 times daily

Ophthalmic Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Dry eye preparations **Vismed** eye drops **10mL**		Each 1mL contains, Sodium hyaluronate 1.8mg	· Corneal conjunctival epithelial disorder due to : - Sjogren's syndrome, Stevens Johnson's syndrome, endogenous diseases such as dry eye syndrome - Exogenous diseases caused by postoperative, drug-induced, trauma, wearing contact lenses, etc. · 1 drop 5~6 times daily
Anti-infectives **Moxikacin** eye drops **5mL**		Each 1mL contains, Moxifloxacin 5mg	· Bacterial conjunctivitis, tarsadenitis, keratitis, keratohelcosis · 1 drop 3 times a day
Anti-infectives **Moxikacin** eye drops **0.4mL**		Each 1mL contains, Moxifloxacin 5mg	· Bacterial conjunctivitis, tarsadenitis, keratitis, keratohelcosis · 1 drop 3 times a day
Anti-infectives **Operan** eye drops **5mL**		Each 1mL contains, Ofloxacin 3mg	· Blepharitis, dacryocystitis, hordeolum, conjunctivitis, keratitis, keratohelcosis, adenophthalmia, infection of ocular surgery · 1 drop 3 times a day
Anti-infectives **Trona** eye drops **5mL**		Each 1mL contains, Tobramycin 3mg	· Blepharitis, dacryocystitis, hordeolum, conjunctivitis, keratitis, keratohelcosis · Mild, moderate infection : 1~2 drops q4h severe infection : 2 drops hourly
Anti-infectives **Operan** Ophthalmic Ointment **5g**		Each 1g contains, Ofloxacin 3mg	· Blepharitis, dacryocystitis, hordeolum, conjunctivitis, keratitis(including keratohelcosis), adenophthalmia, infection of ocular surgery · 1 drop 3 times a day
Anti-infectives **Levokacin** eye drops **5mL**		Each 1mL contains, Levofloxacin 5mg	· Blepharitis, hordeolum, dacryocystitis, conjunctivitis, blepharadenitis, keratitis, corneal ulcer, aseptic therapy in ophthalmic surgery · 1 drop at a time, t.i.d, instillation
Anti-infectives **Levokacin eye** drops **1.5%** **0.4mL**		Each 1mL contains, Levofloxacin 15mg	· Blepharitis, hordeolum, dacryocystitis, conjunctivitis, blepharadenitis, keratitis, corneal ulcer, aseptic therapy in ophthalmic surgery · 1 drop at a time, t.i.d, instillation

Ophthalmic Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Glaucoma preparations **Lanotan** eye drops **2.5mL**		Latanoprost Each 1mL contains. Latanoprost 0.05mg	· Open-angle glaucoma, chronic closed angle glaucoma, ocular hypertension · 1 drop every evening
Glaucoma preparations **Lanope** eye drops **0.2mL**		Latanoprost Each 1mL contains. Latanoprost 0.05mg	· Open-angle glaucoma, chronic closed angle glaucoma, ocular hypertension · 1 drop every evening
Glaucoma preparations **Alphamon-p** eye drops **5mL**		Each 1mL contains Brimonidine tartrate 1.5mg	· Open-angle glaucoma, Ocular hypertension · 1 drop 3 times daily
Glaucoma preparations **Dortisop S** eye drops **0.4mL**		Each 0.4mL contains, Dorzolamide hydrochloride 8.904mg Timolol maleate 2.732mg	· Reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension who are insufficiently responsive to beta-blockers · 1 drop twice a day
Anti-allergy **Allepatadine** eye drops **0.2%, 0.7%** **0.3mL**		Each 1mL contains, Olopatadine HCl	· Treatment of symptoms of allergic conjunctivitis · 1 drop once a day(0.2%, 0.7%)
Anti-allergy **Allepatadine** eye drops **0.1%, 0.2%, 0.7%** **2.5mL, 3mL, 5mL**		Each 1mL contains, Olopatadine HCl 1.11mg, 2.22mg, 7.76mg	· Treat itching in the eyes due to allergic conjunctivitis. · 1 drop 2 times daily(0.1%) · 1 drop once a day(0.2%, 0.7%)
Corticosteroids **Fumetholone** eye drops **5mL**		Each 1mL contains, Fluorometholone 1mg	· Blepharitis, conjunctivitis, keratitis, scleritis, iritis, uveitis, inflammation after surgery, etc. · 1~2 drops 2~4 times daily
Corticosteroids **Fumetholone Plus** **eye drops** **6mL**		Each 1mL contains, Fluorometholone 1mg Tetrahydrozoline 0.25mg	· Noninfectious conjunctivitis, allergic keratitis with severe swelling&congestion. · Noninfectious inflammaion of anterior segment(panuveitis, episcleritis, scleritis) · Post-operative inflammaion · 1 drop 2~3 times daily

Ophthalmic Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Corticosteroids **Fumetholone** eye drops **0.45mL**		Each 1mL contains, Fluorometholone 1mg	· Blepharitis, conjunctivitis, keratitis, scleritis, iritis, uveitis, postoperative inflammation, etc. · 1~2 drops 2~4 times daily
Corticosteroids **Pretesol** eye drops **10mL**		Each 1mL contains, Prednisolone acetate 10mg	· Blepharitis, conjunctivitis, keratitis, sleritis, uveitis, inflammation after surgery · 1 to 2 drops, 2 to 4 times a day
NSAIDs **Eyepecsin** eye drops **5mL**		Each 1mL contains, Diclofenac sodium 1mg	· Prevention of symptoms related cataract surgery : post operative inflammation, miosis during operation, and cystoid macular edema : 1 drop 3~5 times daily(maintenance dose) · Treatment of inflammation in anterior of eye : 1 drop 4~5 times daily
NSAIDs **Broeye** eye drops **0.4mL**		Each 1mL contains, Bromfenac sodium hydrate 1mg	· Symptomatic treatment of inflammatory diseases of the external and anterior segment(blepharitis, conjunctivitis, scleritis(including episcleritis), post-operative inflammation) · 1 drop at a time, b.i.d, instillation
Surgery Supplement **Kukje Hyaluronate** **eye inj.** **0.85mL**		Each 1 syringe(0.85mL) contains, Hyaluronate sodium 12.75mg	· Injection for cataract surgery and intraocular lens impantation, keratoplasty and glaucoma filtering surgery
[OTC] Conjunctivitis & Hordeolum medicines **Synthotop Cool** eye drops **14mL**		Each 1mL contains, Aminocaproic Acid 20mg Chlorpheniramine Maleate 0.2mg Dipotassium Glycyrrhizinate 1mg Sulfamethoxazole 40mg	Keratoplasty and glaucoma filtering surgery
[OTC] Inflamed eye & Eyestrain **Camile Cool** eye drops **14mL**		Each 1mL contains, Aminocaproic Acid 10mg Chlorpheniramine Maleate 0.1mg Naphazoline HCl 0.02mg Neostigmine Methylsulfate 0.02mg	· Eyestrain, conjunctival hyperemia, eye discomfort after swimming or sweating or dust gets in your eyes, when due to the ultraviolet rays, and other eye irritation, fester eyelids, discomfort when wearing hard contact lenses, eye itching, blurry(when a lot of gum, etc.) · 2~3 drops, 5~6 times daily

Ophthalmic Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
[OTC] Ophthalmic Agents **Onceeye** eye drops 1% **0.5mL**		Each 1mL contains, Carboxymethylcellulose sodium 10mg	· For the temporary relief of burning, irritation, and discomfort due to dryness of the eye or exposure to wind or sun · Instill 1 or 2 drops in the affected eye(s) as needed

Cephalosporins

Product name		Ingredient and Content	Indications / Dosage and Administration
1st Generation Cephalosporins **Kukje Cefadroxil** cap.		Each capsule contains, Cefadroxil 500mg(activity)	· Urinary infection, epidermis&skin infection, pharyngitis, respiratory infection, etc. · 1~2g daily in 1~2 divided doses
1st Generation Cephalosporins **Tricef** cap. **250mg, 500mg**		Each capsule contains, Cefradine 250mg, 500mg(activity)	· Tonsillitis, pharyngitis, lobar pneumonia, bronchitis, prostatitis, cystitis, pyelonephritis, etc. · 1~2g daily in 2~4 divided doses
2nd Generation Cephalosporins **Kukje Cefminox** inj. **0.5g, 1g**		Each vial contains, Cefminox sodium 500mg, 1g (activity)	· Cholecystitis, cholangitis, peritonitis, inflammation of pelvic dead space, intrauterine infection, septicemia, peritonsillar abscess, tonsillitis, bronchitis, pneumonia, pyelonephritis, cystitis, etc · 2-4g/day by IV inj/infusion in 2 divided doses, may be increased up to 6 g/day.(adult). 20 mg/kg body wt by IV inj/infusion tid-qid.(children).
1st Generation Cephalosporins **Kukje Cefazedone** inj. **0.5g, 1g, 2g**		Each vial contains, Cefazedone sodium 500mg, 1g, 2g(activity)	· Respiratory infection, urinary tract infection, cholangitis, gynecological infection, skin infection, bone&joint infection, septicemia, endocarditis, surgery infection, etc. · G(+) : 1~2g daily in 2~3 divided doses I.V. G(-) : 3~4g daily in 2~3 divided doses I.V.

Cephalosporins

Product name		Ingredient and Content	Indications / Dosage and Administration
2nd Generation Cephalosporins **Kemocin** dry syr. **Kemocin** cap.		(Dry syr) Each 100g contains, Cefaclor 3.75g(activity) (Cap) Each capsule contains, Cefaclor 250mg(activity)	· Otitis media, pneumonia, laryngopharyngitis, tonsillitis, bronchitis, pyelonephritis, cystitis, gonococcal urethritis, etc. · (Dry syr)0mg/kg daily in 3 divided doses for children · (Cap)250mg q8hr(adult)
2nd Generation Cephalosporins **Kukje Cefotetan** Na. inj. **0.5g, 1g**		Each vial contains, Cefotetan 500mg, 1g(activity)	· Septicemia, superficial 2nd infection, bronchitis, tonsillitis, pyelonephritis, cystitis, cholangitis, peritonitis, intrauterine infection, etc. · 1~2g daily in 2 divided doses I.V. or infusion (adult) 40~60mg/kg daily in 2~3 divided doses I.V. or infusion(children)
2nd Generation Cephalosporins **Zincef** tab.		Each tablet contains, Cefuroxime 250mg(activity)	· Otitis media, sinusitis, bronchitis, pneumonia, pyelonephritis, cystitis, urethritis, furunculosis, gonorrhea, etc. · 250mg b.i.d
3rd Generation Cephalosporins **Kukje Cefixime** cap.		Each capsule contains, Cefixime 100mg(activity)	· Bronchitis, pneumonia, pyelonephritis, cystitis, gonococcal urethritis, cholecystitis, otitis media, scarlet fever, etc. · 50~100mg b.i.d
3rd Generation Cephalosporins **Kukje Ceftriaxone** Na. inj. **0.5g, 1g, 2g**		Each vial contains, Ceftriaxone sodium 500mg, 1g, 2g(activity)	· Respiratory infection, ENT infection, renal&urinary tract infection, septicemia, meningitis, peritonitis, cholecystitis, etc. · 1~2g qd I.V., I.M.(adult), 20~50mg/kg qd(~14 days; infant), 20~80mg/kg qd (15days~12years; children)
3rd Generation Cephalosporins + β Lactamase inhibitor **Peratam** inj. **1g, 2g**		Each vial contains, Cefoperazone sodium 500mg(activity) + Sulbactam sodium 500mg(activity) Cefoperazone sodium 1g(activity) + Sulbactam sodium 1g(activity)	· Bronchitis, pneumonia, pyelonephritis, cystitis, septicemia, cholecystitis, meningitis, pelvic inflammatory disease, endometritis, etc. · 2~4g daily in 2 divided doses I.V.(adult)

Cephalosporins

Product name		Ingredient and Content	Indications / Dosage and Administration
3rd Generation Cephalosporins **Ditoren** Fine granule		Each 1g contains, Cefditoren Pivoxil 100mg	· Urinary tract infection, Laryngopharyngitis, Acute bronchitis, Tonsillitis, Pneumonia, Abscess of lung, Otitis media, Sinusitis, etc. · Pediatrics : 3mg/kg 1 time, t.i.d, PO

Penicillines

Penicillines **Potacin** tab.		Each tablet contains, Amoxicillin hydrate 250mg/ Potassium clavulanate 125mg	· Acute·chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, pelvic infection, gonorrhea, boils and abscesses, cellulitis, wound infection, osteomyelitis, dental infection · Adults and childre over 12 years old or over 40kg in weight : 250mg/125mg once a day, t.i.d(every 8hours), PO (For severe and respiratory infections, the dose may be increased to 500mg/125mg once.)
Penicillines **Potacin duo** tab. **500mg**		Each tablet contains, Amoxicillin hydrate 437.5mg/ Potassium clavulanate 62.5mg	· Acute·chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, pelvic infection, gonorrhea, boils and abscesses, cellulitis, wound infection, osteomyelitis, dental infection · 2 tablets at a time, b.i.d(every 12hours), PO
Penicillines **Potacin** dry syr.		Each 100g(908.15mL) contains, Amoxicillin hydrate 36.3143g (titer)/ potassium clavulanate 5.1878g (titer)	· Acute·chronic bronchitis, lobar and bronchopneumonia, empyema, lung abscess, tonsillitis, sinusitis, otitis media, cystitis, urethritis, pyelonephritis, pelvic infection, gonorrhea, boils and abscesses, cellulitis, wound infection, osteomyelitis, dental infection · Mild to moderate infections : 25mg/3.6mg per kg of body weight per day, b.i.d, PO · Severe infections : 45mg/6.4mg per kg of body weight per day, b.i.d, PO

Quinolones

Product name		Ingredient and Content	Indications / Dosage and Administration
Quinolones **Kipocin** tab. **250mg**		Each tablet contains, Ciprofloxacin 250mg(activity)	· Respiratory infection, ENT infection, urinary tract infection, GI infection, bone and joint infection, septicemia, Ob&Gy infection, infections related eye system, etc. · 250~500mg b.i.d (severe combined infection 750mg b.i.d)
Quinolones **Levokacin** tab. **100mg**		Each tablet contains, Levofloxacin 100mg	· Furunculus, hidradenitis, chronic bronchitis, pyelonephritis, urethritis, enteritis, intrauterine infection, sinusitis, otitis media, cholangitis, etc. · 100mg b.i.d~t.i.d(severe 200mg t.i.d)
Quinolones **Levokacin** tab. **750mg**		Each tablet contains, Levofloxacin 750mg	· Pneumonia(Nosocomial, Community Acquired), SSSI, acute Bacterial Sinusitis, Aucte Bacterial Exacerbation of Chronic Bronchitis, Chronic Bacterial Prostatitis, urinary Tract Infection. · 250~750mg qd
Quinolones **Lebokacin** inj. (Bag) **500mg, 750mg**		Each 1mL contains, Levofloxacin 5mg	· Community-acquired pneumonia, complex urinarytract infections including acute pyelonephritis, complex skin and soft tissue infections, acute sinusitis, acute bacterial exacerbation of chronic bronchitis · 250~500mg qd~b.i.d IV infusion (It can be administered for up to 14 days, and the infusion should take at least 60 minutes.) · In general, administration of this drug should be continued for at least 48-72 hours after the patient has obtained fever-reducing status or evidence of eradication. · In case of severe infectious disease, an increase in the dose should be considered.

Tetracycline

Product name		Ingredient and Content	Indications / Dosage and Administration
For treatment bacterial infections **Kukje Doxycycline cap.** **100mg**		Each capsule contains, Doxycycline hyclate 100mg	· Rickettsial infections, respiratory tract infections, ophthalmic infections, sexually transmitted infections, relapsing fever, plague, tularemia, cholera, Campylobacter fetus infections, brucellosis, bartonellosis, trachoma · When bacteriological testing indicates appropriate susceptibility, for the treatment of infections due to Escherichia coli, Enterobacter aerogenes, Shigella species, Acinetobacter species, Haemophilus influenzae, Klebsiella species · When penicillin is contraindicated, as alternative therapy for yaws, listeriosis, Vincent's infection, actinomycosis, Clostridium species infection · Initial dose : 200mg orally on the first day, given in 2 or 4 divided doses · Maintenance dose: 100 mg orally per day, given once a day or in 2 divided doses · More severe infections: 100mg orally every 12 hours

Macrolides

Macrolides **Roxithrin tab.** **150mg**		Each tablet contains, Roxithromycin 150mg(activity)	· Laryngopharyngitis, otitis media, sinusitis, acute bronchitis, tonsillitis, pneumonia, etc. · 150mg b.i.d

Antifungal & Antiviral Agents

Antifungals **Fucozol cap.** **50mg**		Each capsule contains, Fluconazole 50mg	· Vaginal candidiasis, mucocutaneous candidiasis, candidemia, cryptococcosis, prevention of fungal infections in immunocompromised patients, dermatomycosis, etc. · Cryptococcal infection 400mg followed by 200~400mg once daily for 6~8 wk. systemic candidiasis 400mg followed by 200mg once daily. mucosal candidiasis 50mg once daily for 2 wk. vaginal candidiasis 150mg as a single dose. dermal infections 50mg once daily or 150mg once weekly for 2~4 wk.
Antifungals **Itracon tab.** **100mg**		Each tablet contains, Itraconazole 100mg	· Onychomycosis, dermatophytosis(tinea pedis, tinea corporis, etc), candidial vaginitis, pityriasis versicolor, systemic infections(aspergillosis, cryptococcosis, etc), etc. · Short-term treatment : 100~200mg, once or twice daily · Onychomycosis : 200mg b.i.d for 1week, repeated once for fingernail infections and twice for toenail infections after drug-free intervals of 3weeks, Alternatively, 200mg qd for 3mth. · Systemic infections : 100~200mg qd

Antifungal & Antiviral Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Antiviral Agents **Kukje Acyclovir tab.** **200mg**		Each tablet contains, Acyclovir 200mg	· Herpes simplex virus infection, herpes zoster, varicella · Herpes simplex infections : 200mg 5 times daily every 4 hr for 5 days · Prevention of herpes simplex infections : 200mg 4 times/day · Herpes zoster : 800mg 5 times/day for 7days · Varicella(≥2yrs) : 20mg/kg(max 800mg) 4times/ day for 5 days
Antiviral Agents **Kukje Acyclovir cream** **5g**		Each 1g contains, Acyclovir 50mg	· Herpes simplex virus infection · Apply to the lesion 5 times daily at 4 hours interval

Adrenal Corticosteroids

Corticosteroids **Nisolon M tab.**		Each tablet contatins, Methylprednisolone 4mg	· Adrenal cortical insufficiency, rheumatoid arthritis, systemic lupus erythematosus, multiple sclerosis, severe asthma, stevens-johnson syndrome, iridocyclitis, autoimmune hemolytic anemia, leukemia, lymphoma, etc. · 4~48mg daily (Multiple sclerosis : 200mg/day for a week, followed by 80mg/alternate day treatment for 30 days)

CNS Agents

Antidepressants **Trittico tab.** **25mg, 50mg** **Trittico ER tab.** **75mg**		(tab) Each tablet contains, Trazodone HCl 25mg, 50mg (ER tab) Each tablet contains, Trazodone HCl 75mg	· Depression · 150mg on the first day, increase 50mg every 3~4days, up to 400mg/day(outpatient), 600mg/ day(inpatient)
Antidepressants **Kukje Fluoxetine cap.** **20mg**		Each capsule contains, Fluoxetine 20mg	· Depression, bulimia nervosa, obsessive-compulsive disorder, premenstrual dysphoric disorder · 20mg qd(depression), 60mg/day(bulimia nervosa), 20~60mg(obsessive-compulsive disorder), 20mg/ day(premenstrual dysphoric disorder)
Anticonvulsants **Neopentin cap.** **100mg, 300mg**		Each capsule contains, Gabapentin 100mg, 300mg	· Epilepsy, neuropathic pain · 300mg qd or 100mg t.i.d on first day(300mg/day), 300mg b.i.d or 200mg t.i.d on second day(600mg/ day), 300mg t.i.d from third day(900mg/day; maintenance dose)

CNS Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Cerebrovascular disease **Colensina** soft cap.		Each capsule contains, Choline alphoscerate 400mg	· Memory deterioration, confusion, disorientation disorders due to hypobulia and spontaneity degradation, hypobulia and spontaneity degradation, decreased concentration · Emotional instability, irritability, indifference · Senile pseudo-depression · 400mg 2~3 times daily
Cerebrovascular disease **Nigorine** tab. **30mg**		Each tablet contains, Nicergoline 30mg	· Primary treatment for the following dementia syndromes associated with primary degenerative vascular dementia and mixed dementia: memory impairment, attention deficit, impaired judgment, and lack of initiative. · Adult: 30mg qd, PO, before meals(max 60mg)

Osteoporosis Agents

Osteoporosis Agents **Alendros plus D** tab.		Each tablet contains, Alendronate Na 91.37mg[alendronate 70mg], Cholecalciferol concentrated 53.33mg [Vit D3 5,600IU]	· Osteoporosis in postmenopausal women and men · 1tab./gd/week
Osteoporosis Agents **Ibannate** tab. **150mg**		Each tablet contains, Ibandronate sodium monohydrate 168.75mg	· The treatment and prevention of osteoporosis in postmenopausal women. · 150mg/q.d/month

Antihistamines

Antihistamines **L-trizine** tab.		Each tablet contains, Levocetirizine HCl 5mg	· Allergic rhinitis, chronic idiopathic urticaria, dermatitis and eczema with itching · 5mg qd
Antihistamines **Allepatadine** tab.		Each tablet contains, Olopatadine HCl 5mg	· Allergic rhinitis, Urticaria, skin diseases with itching (eczema, dermatitis, prurigo, pruritus cutaneus, psoriasis vulgaris) · 1tab b.i.d
Antihistamines **Tabeon** tab. **Tabeon** SR tab.		Each tablet contains, Bepotastine besilate 10mg, 20mg	· Multi-annual allergic rhinitis, chronic urticaria, pruritus associated with skin disease(eczema, dermatitis, pruritus cutaneus, prurigo) · 1tab qd

Antihistamines

Product name		Ingredient and Content	Indications / Dosage and Administration
Antihistamines **Epinawon** tab.		Each tablet contains, Epinastine HCl 10mg	1. bronchial asthma 2. Allergic rhinitis 3. Urticaria, eczema·dermatitis, pruritus cutaneus, prurigo, psoriasis with itching · 10~20mg qd

NSAIDs & Other Analgesics

NSAIDs **Arofen** tab.		Each tablet contains, Aceclofenac 100mg	· Rheumatoid arthritis, osteoarthritis, toothache, posttraumatic inflammation, etc. · 1tab b.i.d
NSAIDs **Meloxifen** cap. **7.5mg, 15mg**		Each capsule contains, Meloxicam 7.5mg, 15mg	· Rheumatoid arthritis, osteoarthritis, ankylosing spondylitis · 7.5~15mg qd
NSAIDs **Analoxo** tab.		Each tablet contains, Loxoprofen 60mg	· Rheumatoid arthritis, back pain, cervicoomobrachial syndrome, posttraumatic·postoperative inflammation and pain, upper respiratory inflammation, etc. · 60mg t.i.d, monotherapy : 60~120mg
Analgesics **Hiset** tab. **Hiset** semi tab.		(Hiset tab) Each tablet contains, Acetaminophen 325mg Tramadol HCl 37.5mg (Hiset semi tab) Each tablet contains, Acetaminophen 162.5mg Tramadol HCl 18.75mg	· Moderate-severe acute·chronic pain · (Hiset tab)2tab at more than 6hrs interval (Hiset semi tab)4tab at more than 6hrs interval
Musculoskeletal drugs **Ostera** tab.		Each tablet contains, Angelica gigas Nakai · Chaenomeles sinensis (Thouin) Koehne · Saposhnikovia divaricata Schischkin ·Dipsacus asperoides C.Y.Cheng et T.M.Ai · Acanthopanax sessiliflorum Seeman · Achyranthes japonica Nakai · Clematis chinensis Osbeck · Cinnamomum cassia Presl · Gentiana macrophylla Pallas · Cnidium officinale Makino · Gastrodia elata Blume · Carthamus tinctorius Linne 25% Ethanol ext. (3.5→1) 405.4mg	· Relieve symptoms of osteoarthrosis · 1tab b.i.d, PO

NSAIDs & Other Analgesics

Product name		Ingredient and Content	Indications / Dosage and Administration
NSAIDs **Celexib** cap. **200mg**		Each capsule contains, celecoxib 200mg	· Rheumatoid arthritis, osteoarthritis, ankylosing spondylitis, acute pain, primary dysmenorrhea · 1capsule qd
Analgesics & Antipyretics **Amiset** inj.		Each 1ml contains, Acetaminophen 10mg	· Treat moderate to severe pain · Reduce fever Parenteral : - Weight 50kg or greater : 1000mg IV Maximum Single Dose : 1000mg Minimum Dosing Interval : every 4 hours Maximum Dose : 4000mg per 24 hours - 33 to 49Kg : 15mg/kg IV Maximum Single Dose : 15mg/kg Minimum Dosing Interval : every 4 hours Maximum Dose : 60mg/kg per 24 hours or 4000mg per 24 hours · Infuse IV over 15 minutes

Gastrointestinal Agents

Antiulcerant ; H2 blockers **Kukje Cimetidine** tab. **200mg**		Each tablet contains, Cimetidine 200mg	· Gastroduodenal ulcer, GERD, zollinger-Ellison syndrome, gastritis, etc. · 400mg b.i.d or 800mg qd or 300mg q.i.d(adult)
Antiulcerant ; H2 blockers **Axadin** cap. **150mg**		Each capsule contains, Nizatidine 150mg	· Gastroduodenal ulcer, GERD, gastritis, etc. · 150~300mg qd or b.i.d
Antiulcerant ; PPIs **Lanster** cap. **15mg, 30mg**		Each capsule contains, Lansoprazole 15mg, 30mg	· Gastric ulcer, duodenal ulcer, GERD, H. pylorieradication, esophagitis, etc. · 15~30mg qd, H. pylori eradication : 30mg 2~3times daily with antibiotics
Antiulcerant **Kukje Rebamipide** tab. **100mg**		Each tablet contains, Rebamipide 100mg	· Gastric ulcer, gastritis · 100mg t.i.d
GI Tract Regulators **Tarabutin** tab. **100mg, 200mg**		Each tablet contains, Trimebutine maleate 100mg, 200mg	· Esophageal reflux and hiatus hernia, gastroduodenitis, and digestive dysfunction (abdominal pain, dyspepsia, nausea, vomiting) in gastroduodenal ulcer · Irritable bowel syndrome and spastic colon · Pediatric diseases: habitual vomiting, non-infectious intestinal transit disorder (constipation, diarrhea) · 100-200mg t.i.d, PO, before meals

Gastrointestinal Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
GI Tract Regulators **Kukje Itoride** tab.		Each tablet contains, Itopride HCl 50mg	· The relief of the symptoms of functional dyspepsia(anorexia, epigastralgia, sense of abdominal fullness, chest pain, nausea, vomiting · Itopride 50 mg three times daily, before meals.
Antiulcerant ; PPIs **Nexipra** tab. **20mg, 40mg**		Each tablet contatins, Esomeprazole 20mg, 40mg	· GERD, risk reduction of NSAID-associated gastric ulcer, etc. 20~40mg qd · H.pyrori eradication 20mg bid with antibiotics
Antiulcerants **Algitect** sol.		Each suspension contains, Alginate sodium 1g/20mL/pack	· Treatment of gastric ulcers; reflux and pepticesophagitis; treatment of NSAID- or aspirin-induced · GI symptoms and mucosal damage; prevention of stress ulcers and GI bleeding in critically ill patients. · 1g orally on an empty stomach 4 times a day or alternatively, 2g orally on an empty stomach 2 times a day
Antiulcerants **Stil U 2X** tab.		Each tablet contains, Artemisia asiatica 95% ethanol extract(20→1) 90mg	· Improve gastric mucosal lesion(erosion, bleeding, flare,edema) of acute and chronic gastritis · 1tab, b.i.d, PO
GI Tract Regulators **Kukje Mosapride SR** tab.		Each tablet contains, Mosapride citrate anhydrate 15mg	· The relief of the symptoms of functional dyspepsia(Epigastric pain, Nausea, Vomiting) · 1tab QD PO
Antiulcerants; H_2 blockers **Nafudi** tab.		Each tablet contains, Lafutidine 10mg	· Gastric ulcer, duodenal ulcer, improving gastric mucosal lesions of acute gastritis and chronic gastritis, antibiotic combination therapy for pepticulcer patients infected with Helicobacterpylori, pre-anesthesia medication, treatment of reflux esophagitis · Gastric ulcer, duodenal ulcer, treatment of reflux esophagitis : 10mg b.i.d · Improving gastric mucosal lesions of acute gastritis and chronic gastritis : 10mg HS
Antiulcerants **Still U** tab.		Each tablet contains, Artemisia asiatica 95% ethanol extract.(20->1) 60mg (as Eupatilin 0.48-1.44mg and Jaceosidin 0.15-0.45mg)	· Improving gastric mucosal lesions (erosion, bleeding, redness, swelling) of the following diseases : acute gastritis, chronic gastritis, pevention of gastritis by administering NSAID · 3 times daily
Antispasmodic **Propa** tab.		Each tablet contains, Tiropramide hydrochloride 100mg	· Acute spasmodic pain in the following diseases : hepatic and biliary pain, abdominal pain due to various causes, kidney and ureteral pain · Abdominal spasms and pain in the following diseases : gastrointestinal dyskinesia, cholelithiasis, cholecystitis, postoperative adhesion · 100mg 2~3 time daily

Cardiovascular Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Antihyperlipidemic **Apecor tab.**		Each tablet contains, Atorvastatin calcium 10.36mg (as atrovastatin 10mg) Atorvastatin calcium 20.72mg (as atorvastatin 20mg)	· Dyslipidemia · The usual starting dose is 10mg once a day the maximum dose is 80mg once a day.
Antihyperlipidemic **Crerova tab.** **5mg, 10mg, 20mg**		Each tablet contains, Rosuvastatin calcium 5.2mg (as rosuvastatin 5mg) Rosuvastatin calcium 10.4mg (as rosuvastatin 10mg) Rosuvastatin calcium 20.8mg (as rosuvastatin 20mg)	· Dyslipidemia · The usual starting dose is 5mg once a day. the maximum dose is 20mg once a day.
Antihyperlipidemic **Ezerova tab.** **10/5mg, 10/10mg, 10/20mg**		Each tablet contains, Ezetimibe 10mg/ Rosuvastatin calcium 5.2mg (Rosuvastatin 5mg), 10.4mg (Rosuvastatin 10mg), 20.8mg (Rosuvastatin 20mg)	· Primary hypercholesterolemia · 1tab QD
Antihyperlipidemic **Apezet tab.** **10/10mg**		Each tablet contains, Ezetimibe10mg/ Atorvastatin10mg	· Primary Hypercholesterlemia · 1tab Q.D
Antithrombotics **Plagrel tab.** **75mg**		Each tablet contains, Clopidogrel 75mg	· Acute coronary syndrome(ACS), recent MI, recent stroke, or established peripheral arterial disease · 75mg qd
Antithrombotics **Ciletal SR cap.** **200mg**		Each capsule contains, Cilostazol 200mg	· Improvement of ischemic symptoms such as ulcers, pain, and coldness due to chronic occlusive arterial disease (Buerger's disease, Arteriosclerosis obliterans, Diabetic polyneuropathy, etc.) · Inhibition of recurrence after development of cerebral infarction (excluding cardiogenic brain embolism) · 1tab Q.D
Antithrombotics **Antigrel SR tab.**		Each tablet contains, Sarpogrelate hydrochloride 300mg	· Improvement of ischemic preparation symptoms such as ulcers, pain, and coldness due to chronic occlusive arterial disease (Buerger's disease, Arteriosclerosis obliterans, Diabetic polyneuropathy, etc.) · 1tab Q.D after meals

Cardiovascular Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Antihypertensives; ARB **Kukje Losartan tab.** **50mg**		Each tablet contains, Losartan potassium 50mg	· Hypertension, diabetic nephropathy in Type2 DM(with hypertension) · 50~100mg qd
Antihypertensives; ARB+Diuretics **Kukje Losartan plus tab..**		(Kukje losartan plus tab) Each tablet contains, Losartan potassium 50mg Hydrochlorothiazide 12.5mg	· Hypertension · Losartan potassium/hydrochlorothiazide 50/12.5mg~100/25mg qd
Antihypertensives; CCB **Kukje Amlodipine tab.** **5mg**		Each tablet contains, Amlodipine 5mg	· Hypertension, stable angina, cardiac ischemia · 5~10mg qd
Antihypertensives; β-Blocker **Vasodilren tab.**		Each tablet contains, Carvedilol 25mg	· Hypertension, chronic stable angina · Hypertension : 12.5~25mg qd(max 25mg b.i.d) chronic stable angina : 12.5~25mg b.i.d (max 50mg b.i.d)
Antihypertensives : ARB **Valsar tab.** **80mg, 160mg**		Each tablet contains, Valsartan 80mg Valsartan 160mg	· Hypertension, congestive heart failure · Hypertension : 80mg qd congestive heart failure : 40~160mg qd
Antihypertensives : ARB+CCB **Exduo tab.** **5/80mg, 5/160mg, 10/160mg**		Each tablet contains, - 5/80mg : Amlodipine besylate 6.94mg (as amlodipine 5mg) + Valsartan 80mg - 5/160mg : Amlodipine besylate 6.94mg (as amlodipine 5mg) + Valsartan 160mg - 10/160mg : Amlodipine besylate 13.87mg (as amlodipine 10mg) + Valsartan 160mg	· Hypertension · 1tab qd
Other Antihypertensives **Dualstar tab.** **40/5mg, 40/10mg, 80/5mg**		Each tablet contains, Telmisartan 40mg, 80mg/Amlodipine besylate 6.935mg(Amlodipine 5mg), 13.87mg(Amlodipine 10mg)	· Hypertension which of telmisartan or amlodipine monotherapy can not adequately control blood pressure · 1tab QD
Antilipidemic agents **Omecin soft cap.**		Each capsule contains, Omega-3-acid ethyle esters 90 1000mg	· Dietary supplement for reducing increased triglyceride levels in endinent high triclyceride blood patient · 2g~4g daily, 1~2times

Antidiabetics

Product name		Ingredient and Content	Indications / Dosage and Administration
Antidiabetics **Dymeryl** tab.		Each tablet contains, glimepiride 2mg	· Type 2 diabetes mellitus · 1tab qd
Antidiabetics **Glavis** XR tab. **500mg**		Each tablet contains, Metformin HCl 500mg	· Treatment of adult patients with type 2 diabetes mellitus with insufficient glycemic control through diet and exercise. (especially those with diabetes who are overweight) · This drug can be used as monotherapy or in combination with other oral hypoglyemic agents or insulin. · 500mg Q.D, PO, with dinner. (max 2000mg/ daily) · When increasing the dose, it should be increased by 500mg every week. If blood sugar control is not achieved even after using up to 2000mg once a day, consider administering 1000mg b.i.d, and both times should be administered with meals.
Antidiabetics **Glavis** tab. **1000mg**		Each tablet contains, Metformin hydrochloride 1000mg	· Treatment of Type 2 Diabetics with insufficient blood glucose control through Diet and Exercise Therapy(In particular, overweight diabetics) In adults, it can be used alone or in combination with other oral hypoglycemic agents or insulin. In children 10 years of age and older in adolescents, the drug can be used alone or in combination with insulin. · 500mg b.i.d or t.i.d, PO, with a meal(Dosing must be started at a low dose.) The dose increase is gradually increased by 500mg weekly, and when administered up to 2000mg is usually divided into twice daily morning and evening. If it exceeds 2000mg, it can be divided into three meals per day with meals and up to 2550mg per day.
Antidiabetics **Kukje Pioglitazone** tab. **15mg, 30mg**		Each tablet contains, Pioglitazone HCl 16.53mg (as Pioglitazone 15mg) Pioglitazone HCl 16.53mg (as Pioglitazone 30mg)	· Type 2 diabetes mellitus · Monotherapy : 15mg~30mg QD · Combination therapy(sulfonylurea or metformin or insulin) : 15mg QD
Antidiabetics **Piovis** tab.		Each tablet contains, Pioglitazone HCl 16.53mg (as Pioglitazone 15mg) Metformin hydrochloride 850mg	· Type 2 diabetes mellitus · 1tab QD(max Pioglitazone 30mg/Metformin 2500mg)
Antidiabetics **R-dect** tab. **480mg**		Each tablet contains, R-thioctic acid thomethamine 480mg (as R-thiotic acid 300mg)	· Alleviation of diabetic multiple neuritis · 480mg QD


Antidiabetics

Product name		Ingredient and Content	Indications / Dosage and Administration
Antidiabetics **Teneldi** tab. **20mg**		Each tablet contains, Teneligliptin hydrochloride hydrate 26mg (as Teneligliptin 20mg)	· It is administered as an adjunct to diet and exercise therapy to improve blood sugar control in adults with type 2 diabetes mellitus. · It is administered as a monotherapy. · It is administered as a comination therapy in the following cases. (When metformin monotherapy cannot adeuately control blood sugar / When metformin and sulfonylurea combination therapy cannot achieve adequate blood sugar control) · 20mg Q.D, PO
Antidiabetics **Teneldi-M** SR tab. **10/500mg, 10/750mg, 20/1000mg**		Each tablet contains, Teneligliptin Hydrochloride hydrate 26mg (as Teneligliptin 20mg)/ Metformin HCl 1000mg, Teneligliptin Hydrochloride hydrate 13mg(as Teneligliptin 10mg)/ Metformin HCl 750mg, 500mg	· It is administered as an adjunct to diet and exercise therapy to improve blood sugar control in type 2 diabetes patients, for which the combination of the two components is suitable. · For 20/1000mg, 1tab, Q.D, PO · For 10/750mg or 10/500mg, 2tabs, Q.D, PO · The maximum recommended daily dose is teneligliptin 20mg and SR metformin 2000mg.
Antidiabetics **Forxidi** tab. **10mg**		Each tablet contains, Dapagliflozin 10mg	· Type 2 diabetes melitus · 1tab, Q.D
Antidiabetics **Forxidi-M** XR tab. **10/500mg, 10/1000mg**		Each tablet contains, Dapagliflozin 10mg/ Metformin HCl 500mg Dapagliflozin 10mg/ Metformin HCl 1000mg,	· Type 2 diabetes melitus · Generally Q.D with dinner
Antidiabetics **Forxivis** tab. **10/100mg**		Each tablet contains, Dapagliflozin 10mg/ Sitagliptin 100mg	· Type 2 diabetes melitus · 1tab qd
Antidiabetics **Janudi** tab. **25mg, 50mg, 100mg**		Each tablet contains, Sitagliptin 25mg, 50mg, 100mg	· Type 2 diabetes melitus - Monotherapy - Combination therapy · 1tab qd
Antidiabetics **Gliance** tab. **10mg, 25mg**		Each tablet contains 10 mg or 25 mg of empagliflozin	· An adjunct to diet and exercise to improve glycemic control in patients with type 2 diabetes mellitus · 10–25 mg once daily
Antidiabetics **Gliance Duo** tab. **5/500mg, 5/850mg, 5/1000mg, 12.5/500mg, 12.5/850mg, 12.5/1000mg**		Each tablet contains 5 mg or 12.5 mg of empagliflozin and 500 mg, 850 mg, or 1000 mg of metformin hydrochloride	· An adjunct to diet and exercise to improve glycemic control in patients with type 2 diabetes mellitus for whom combination therapy with empagliflozin and metformin is appropriate · One tablet twice daily maximum recommended daily dose: empagliflozin 25 mg, metformin 2000 mg.

Antidiabetics

Product name		Ingredient and Content	Indications / Dosage and Administration
Antidiabetics **Janudi-M tab.** **50/500mg, 50/850mg, 50/1000mg**		Each tablet contains, Sitagliptin 50mg/ Metformin HCl 500mg, Sitagliptin 50mg/ Metformin HCl 850mg, Sitagliptin 50mg/ Metformin HCl 1000mg	· Type 2 diabetes melitus · The maximum recommended daily dose is Sitagliptin 100mg and Metformin 2000mg. · 1tab b.i.d
Antidiabetics **Janudi-M XR tab.** **50/500mg, 50/1000mg, 100/1000mg**		Each tablet contains, Sitagliptin 50mg/ Metformin HCl 500mg, Sitagliptin 50mg/ Metformin HCl 1000mg, Sitagliptin 100mg/ Metformin HCl 1000mg	· Type 2 diabetes melitus · The maximum recommended daily dose is Sitagliptin 100mg and Metformin 2000mg. · 100/1000mg 1tab qd · 50/500mg, 50/1000mg 2tab qd
Antidiabetics **Tradi-M tab.** **5mg**		Each tablet contains, Linagliptin 5mg	· Type 2 diabetes melitus - Monotherapy - Combination therapy · 1tab qd
Antidiabetics **Tradi-M Duo tab.** **2.5/500mg, 2.5/850mg, 2.5/1000mg**		Each tablet contains, Linagliptin 2.5mg/ Metformin HCl 500mg Linagliptin 2.5mg/ Metformin HCl 850mg Linagliptin 2.5mg/ Metformin HCl 1000mg	· Type 2 diabetes melitus · The maximum recommended daily dose is Linagliptin 5mg and Metformin 2000mg. · 1tab b.i.d

Genitourinary Agents

Drugs for Prostate Disorder **Utanal cap.**		Each capsule contains, Tamsulosin HCl 0.2mg	· Benign prostatic hyperplasia · 0.2mg qd
Drugs for Prostate Disorder **Utanal SR cap.** **0.4mg**		Each capsule contains, Tamsulosin HCl 0.4mg	· Benign prostatic hyperplasia · 1tab, Q.D, after meals
Drugs for Prostate Disorder **Finastid tab.**		Each tablet contains, Finasteride 5mg	· Benign prostatic hyperplasia · 5mg qd
BPH 5α-Reductase inhibitor (*BPH:benign prostatic hyperplasia) **Avosted soft cap.**		Each soft capsule contains, Dutasteride 0.5mg	· Benign prostatic hyperplasia, acute urinary retention, androgenetic alopecia · 1capsule qd
Erectile dysfunction medicines : PDE 5 inhibitor **Victoris tab.** **5mg, 20mg**		Each tablet contains, Tadalafil 5mg, 20mg	· Erectile dusfunction · 10mg qd(recommended), increase to 20mg qd based on efficacy and tolerability.

Genitourinary Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Erectile dysfunction medicines : PDE 5 inhibitor **Maxigra tab.** **100mg**		Each tablet contains, Sildenafil citrate 140.45mg (as sildenafil 100mg)	· Treatment of erectile dysfunction · 25~50mg approximately 1hour before sexsual activity
Erectile dysfunction medicines : PDE 5 inhibitor **Maxigra ODT·** **50mg** **(*ODT : orally disintegrating tablet)**		Each tablet contains, Sildenafil citrate 70.23mg (Sildenafil 50mg)	· Erectile dusfunction · 25~50mg 50mg qd prn 1 hr(recommended) or anywhere from 30 min to 4 hrs before sexsual activity
Trazosin tab.		Each tablet contains 5.935 mg of terazosin hydrochloride hydrate (equivalent to 5 mg of terazosin)	·Urinary disorders due to benign prostatic hyperplasia, hypertension (mild to moderate) 1.Urinary disorders due to benign prostatic hyperplasia: Initial dose of 1 mg orally, maintenance dose of 5–10 mg orally once daily. 2.Hypertension: Initial dose of 1 mg orally, maintenance dose of 2–10 mg orally once daily.

Respiratory Agents

Antitussives & Expectorants **Erpect cap.**		Each capsule contains, Erdosteine 300mg	· Mucolytics and expectorants for chronic and acute respiratory disease · 1cap 2~3 times daily
Muscle Relaxants **Ericson tab.**		Each tablet contains, Eperisone HCl 50mg	· Muscle relaxant that has been used in the symptomatic treatment of muscle spasm and spasticity(cervicoomobrachial syndrome, low back pain, etc) · 50mg t.i.d
Skeletal muscle relaxants **Ericsone SR tab.**		Each tablet contains, Eperisone hydrochloride 75mg	· Painful muscle spasms with musculoskeletal disorders : Cervicoomobrachial syndrome, Frozen shoulder, Backache · 1tab b.i.d, PO, after meals · Since it is a slow release tablet, take it whole without crushing, splitting, or chewing.
Vasodilators **Kukje Limaprost tab.**		Each tablet contains, Limaprost 5μg	· Buerger's disease, acquired stenosis · Buerger's disease - 10μg/t.i.d/day(30μg/day) acquired stenosis - 5μg/t.i.d/day(15μg/day)
Chronic hepatitis **Pafecton Plus cap.**		Each capsule contains, Biphenyl dimethyl Dicarboxylate 12.5mg Ursodeoxycholic acid 50mg	· Chronic persistent hepatitis · 1cap t.i.d
Pre-filled Syringe for Gonarthrosis **Kukje Hyal inj.**		Each 1mL contains, Sodium Hyaluronate 10mg	· Gonarthrosis · Periarthritis of shoulder · Administer 25mg once a week for 5 consecutive weeks
Pre-filled Syringe for Osteoarthritis of Knee Joint **Hyaluronic Acid Joint inj.**		Each 1mL contains, Sodium hyaluronate 10mg	· Osteoarthritis of the knee joint · Treatment of Periarthritis of shoulder · Usually administered once a week to an adult for three consecutive weeks

Respiratory Agents

Product name		Ingredient and Content	Indications / Dosage and Administration
Anti-allergic thinitis **Nasocure** nasal spray		Each 1mL contains, Mometasone furoate 0.5mg	· Adults and children over 2 years old : Seasonal allergic rhinitis and year-round rhinitis Adults over 18 years old : Nasal polyp Adults and youth over 12 years old : Acute rhinosinusitis without severe bacterial infection · Allergic rhinitis - Adults (including the elderly) and youth over 12 years old : In general, spray 2 times in each nostril once a day. (Total dose 0.2mg per day). - Children aged 2 to 11 : In general, spray once a day, once for each nostril. (Total dose 0.1mg per day). · Nasal polyp : In general, spray once a day, twice for each nostril. (Total dose 0.2mg per day). · Acute Rhinosinusitis : In general, spray 2 times a day, twice for each nostril.(Total dose 0.4mg per day).

Other ETCs

Asthma & Allergic rhinitis **Onceair** tab. **10mg**		Each tablet contains, Montelukast sodium 10mg	· Prevention and continuous treatment of asthma · Mitigate seasonal and year-round allergic rhinitis symptoms · (Audult and youth over 15 years of age) 1tab Q.D · For asthmatic patients, take it in the evening · In the case of allergic rhinitis patients, the medication time is determined according to the patient's condition · Patients with both asthma and allergic rhinitis take 1tab Q.D and 1tab at a time in the evening

Health Functional Food

Health Functional Food **Femi Balance Plus**		Respecta®(Lactobacillus acidophilus GLA-14™, Lactoferrin Extract, Lactobacillus rhamnosus HN001™)	· It can suppress lactobacillus growth and harmful bacteria and help facilitate bowel movements · It can help to increase beneficial bacteria in the vagina and suppress harmful bacteria
Health Functional Food **Ocutein3**		4 capsules contains Lutein 10mg, Zeaxanthin 2mg, Omega-3(EPA+DHA) 1,000mg, Vit C 500mg, Vit E, 4001U(269mg a-TE), Zinc 12mg, Copper 2mg	· Lutein & Zeaxanthin : To maintain macular pigment density(filter blue light) · Omega-3 : To improve dry eye syndrome, blood circulation and retinal constituent · Anti-oxidative nutrients(Vit C, Vit E, Zinc, Copper) : To protect visual acuity and provide nutrient to eyes · Take 2 soft capsules 2 times a day with a full glass of water